



Cow-calf Vaccination Programs

Modified Live Virus (MLV) Vaccine Protocol

Open Cows - Vaccinated 4 to 8 weeks prior to breeding

Express[®] FP 5-VL5
Alpha[™] -7/MB-1 or OCU-Guard[®] MB-1
Choice of one of the following dewormers:
LongRange[®] Injection (Rx)
Eprinex[®] Pour-on
IVOMEC[®] 1% Injection
IVOMEC[®] Plus Injection
IVOMEC[®] Pour-on

Calves 2 to 3 Months of Age – Vaccinated at same time open cows are given pre-breeding vaccines

Pyramid 5 + Presponse SQ
Alpha[™] -7/MB-1
Choice of one of the following dewormers:
LongRange[®] Injection (Rx) or
Eprinex[®] Pour-on or
IVOMEC[®] 1% Injection or
IVOMEC[®] Plus Injection or
IVOMEC[®] Pour-on

Killed Virus Vaccine Protocol

Bred Cows

Triangle 10 HB or Triangle 5 and Citadel[™] VL5
Alpha[™] -7/MB-1 or OCU-Guard[®] MB-1
Choice of one of the following dewormers:
LongRange[®] Injection (Rx) or
Eprinex[®] Pour-on or
IVOMEC[®] 1% Injection or
IVOMEC[®] Plus Injection or
IVOMEC[®] Pour-on

Calves 2 to 3 Months of Age

Triangle 5
Presponse HM
Alpha[™] -7/MB-1
Choice of one of the following dewormers:
LongRange[®] Injection (Rx) for calves over 3 months or
Eprinex[®] Pour-on or
IVOMEC[®] 1% Injection or
IVOMEC[®] Plus Injection or
IVOMEC[®] Pour-on