



2023 Tennessee Beef Alliance Health Protocol



Select One Brand, One Box, & an Approved De-wormer for a complete protocol

All HEIFERS are **REQUIRED** to have a shot of Lutalyse or Synchsure on **second working**
(Consult your Veterinarian for this product) ****Administer IM per label Directions****



Pre-Weaning:

1st Working (2 weeks prior to weaning)

- TRIANGLE 5
- Presponse HM
- Bar-Vac 7/Somnus

Use Approved De-wormer

2nd Working (14-28 days later) (Following Weaning)

- EXPRESS 5 or Pyramid 5
- Bar-Vac 7/Somnus

Pre-Weaning:

1st Working (2 weeks prior to weaning)

- Cattlemaster GOLD FP 5
- Ultrabac 7 Somubac

Use Approved De-wormer

2nd Working (14-28 days later) (Following Weaning)

- Bovi-Shield GOLD ONE SHOT (Combo)
- Ultrabac 7 Somubac

Post or at Weaning:

1st Working

- EXPRESS 5 HS
- Presponse HM

Use Approved De-wormer

2nd Working (14-28 days later)

- Express 5 HS
- Alpha-7 or Alpha-7/MB-1

Post or at Weaning:

1st Working

- Bovi-Shield GOLD ONE SHOT (combo)
- Ultrabac 7 Somubac

Use Approved De-wormer

2nd Working (14-28 days later)

- Bovi-Shield GOLD 5
- Ultrabac 7 Somubac

Post or at Weaning:

1st Working

- Pyramid 5 + Presponse SQ (Combo)
- Bar-Vac 7/Somnus

Use Approved De-wormer

2nd Working (14-28 days later)

- Pyramid 5
- Bar-Vac 7/Somnus

Post or at Weaning

1st Working:

- Inforce 3
- One Shot BVD
- Ultrabac 7 Somubac

Use Approved De-wormer

2nd Working (14-28 days later)

- Bovi-Shield GOLD 5
- Ultrabac 7 Somubac

Approved De-wormers Dectomax Injectable or Pour-on, Eprinex/Ivomec Inj. or Pour On/LongRange

De-wormers listed above are approved for any box - NO GENERIC DE-WORMERS

For More Information on Health Protocols contact:

Dr. Locklar-Vet Advisor
931-388-4750

Dr. Boyd-Vet Advisor
931-528-8805

Dr. Malone-Vet Advisor
931-638-1458

Todd Jennings – Boehringer
931-842-0450

Lee Myers - Zoetis
615-934-2486

PLEASE MAINTAIN ALL SERIAL AND LOT NUMBERS FROM VACCINES FOR HEALTH RECORDS

*Note if over 100 days past last shot, another round of the combo vaccines, blackleg, and Lutalyse or Synchsure must be given. * 2nd shot completed no later than 2 weeks prior to sale, or more than 100 days prior to sale.